WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6: C07C 41/14, 43/12

(11) International Publication Number:

WO 97/25303

A1

(43) International Publication Date:

17 July 1997 (17.07.97)

(21) International Application Number:

PCT/GB96/03001

(22) International Filing Date:

6 December 1996 (06.12.96)

(30) Priority Data:

9600072.4

4 January 1996 (04.01.96)

GB

(71) Applicant (for all designated States except US): IMPERIAL CHEMICAL INDUSTRIES PLC [GB/GB]; Imperial Chemical House, Millbank, London SW1P 3JF (GB).

(72) Inventors; and

- (75) Inventors/Applicants (for US only): RYAN, Thomas, Anthony [GB/GB]; "Udine", Quarry Lane, Kelsall, Cheshire CW6 OPD (GB). BURGESS, Leslie [GB/GB]; I Poolside Road, Runcom WA7 5QQ (GB).
- (74) Agents: GEARY, Stephen et al.; ICI Chemicals & Polymers Limited, Intellectual Property Dept., P.O. Box 11, The Heath, Runcorn, Cheshire WA7 4QE (GB).

(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, Cl, CM, GA, GN, ML, MR, NE, SN, TD, TG).

Published

With international search report.

(54) Title: PROCESS FOR THE PRODUCTION OF FLUOROMETHYLHEXAFLUOROISOPROPYLETHER

(57) Abstract

Process for producing Sevoflurane anaesthetic which comprises reacting hexafluoroisopropyl alcohol with essentially pure bis(fluoromethyl)ether. The bis(fluoromethyl)ether is preferably obtained by the reaction of formaldehyde with hydrogen fluoride.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
вв	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	1T	Italy	PL	Poland
BJ	Benin	JР	Japan	PT	Portugal
BR	Brazil	· KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgystan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic	\$D	Sudan
CF	Central African Republic		of Korea	SE	Sweden
CG	Congo	KR	Republic of Korea	SG	Singapore
СН	Switzerland	KZ	Kazakhstan	SI	Słovenia
CI	Côte d'Ivoire	Ll	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LR	Liberia	SZ	Swaziland
CS	Czechoslovakia	LT	Lithuania	TD	Chad
CZ	Czech Republic	LU	Luxembourg	TG	Togo
DE	Germany	LY	Latvia	TJ	Tajikistan
DK	Denmark	MC	Monaco	TT	Trinidad and Tobago
EE	Estonia	MD	Republic of Moldova	UA	Ukraine
ES	Spain	MG	Madagascar	UG	Uganda
FI	Finland	ML	Mali	us	United States of America
FR	France	MN	Mongolia	uz	Uzbekistan
GA	Gabon	MR	Mauritania	VN	Viet Nam

PROCESS FOR THE PRODUCTION OF FLUOROMETHYLHEXAFLUOROISOPROPYLETHER.

5

10

15

20

25

30

This invention relates to a process for the production of fluoromethylhexafluoroisopropylether of formula CH₂F.CH(CF₃)₂ which is the anaesthetic "Sevoflurane".

Several processes have been proposed for the production of Sevoslurane including the reaction of hexasluoroisopropyl alcohol, (CF₃)₂CHOH, with formaldehyde and hydrogen fluoride. One such process comprising adding hexasluoroisopropyl alcohol to a mixture of paraformaldehyde and hydrogen fluoride plus sufficient sulphuric acid to sequester most of the water formed at a temperature above 57°C is described in US 4,250,334. A similar process comprising mixing hexasluoroisopropyl alcohol, formaldehyde, hydrogen fluoride and a dehydrating agent such as sulphuric acid is described in US 4,469,898.

A process for producing an alpha-fluoroether such as and including Sevoflurane is described in International Patent Publication No WO 93/12057, the process comprising reacting a non-enolisable aldehyde such as formaldehyde with hydrogen fluoride to form an intermediate and reacting the intermediate with an alcohol such as hexafluoroisopropyl alcohol to form an alpha-fluoroether such as Sevofluorane. The production of Sevoflurane by adding hexafluoroisopropyl alcohol to the reaction mixture derived from trioxane (as the source of formaldehyde) and hydrogen fluoride and containing the intermediate bis(fluoromethyl)ether is described in Example 19

As is described in WO 93/12057, the reaction products obtained in Example 19 comprised mainly unreacted hexafluoroisopropyl alcohol (72% by mass spectroscopic analysis) and unreacted bis(fluoromethyl)ether (22%) and the yield of Sevoflurane was only 4.9%. This very low yield of Sevoflurane renders the process unsuitable or at best barely suitable for industrial application even with recovery and recycle of unreacted hexafluoroisopropyl alcohol and bis(fluoromethyl)ether.

The present invention is based on the discovery that separation of the intermediate bis(fluoromethyl)ether from the reaction mixture before addition of the hexafluoroisopropyl alcohol results in a process in which Sevoflurane is obtained in high yield and in particular can be obtained as the major product of the reaction.

5

10

15

20

25

30

According to the present invention there is provided a process for the production of fluoromethylhexafluoroisopropylether [Sevoflurane] which comprises reacting essentially pure bis(fluoromethyl) ether with hexafluoroisopropyl alcohol and recovering the resulting fluoromethylhexafluoroisopropylether from the reaction products.

The bis(fluoromethyl)ether is conveniently and preferably produced by the reaction of formaldehyde or a source of formaldehyde with hydrogen fluoride and according to a particular embodiment of the invention there is provided a process for the production of fluoromethylhexafluoroisopropylether which comprises reacting formaldehyde with hydrogen fluoride to produce a reaction mixture containing bis(fluoromethyl)ether, separating essentially pure bis(fluoromethyl)ether from the reaction mixture, reacting the resulting essentially pure bis(fluoromethyl)ether with hexafluoroisopropyl alcohol and recovering the resulting fluoromethylhexafluoroisopropylether from the reaction products.

The reaction between the bis(fluoromethyl)ether and the hexafluoroisopropyl alcohol is conveniently carried out at ambient temperature, say 20°C to 30°C and at atmospheric pressure, although if desired subatmospheric or superatmospheric pressure and a range of temperatures from about 0°C to about 100°C may be employed. The reaction is preferably carried out in the presence of an acid such as sulphuric acid. A slight exotherm resulting in a rise in temperature of the reaction mixture may be observed but in general there is no real advantage in applying heat to the reaction mixture.

The reaction can be readily carried out to result in complete conversion of the hexafluoroisopropyl alcohol and with an acceptable selectivity to the desired Sevoflurane product although a significant amount, say 20% or more, of the acetal (CF₃)₂CHOCH₂OCH₂F is usually produced as a by-product. It is a matter of mere routine experimentation to optimise the reaction conditions to maximise the yield of Sevoflurane and minimise the yield of by-products such as the acetal. A reaction product mixture in which Sevoflurane is the major component with a yield of at least about 55% and preferably about 60% or more is obtained. Carrying out the reaction with complete conversion of the hexafluoroisopropyl alcohol obviates the need to recover and recycle unreacted alcohol.

The amounts of bis(fluoromethyl)ether and hexafluoroisopropyl alcohol will usually be such that the molar ratio of the ether to the alcohol is from about 0.5: 1 to about 1.5: 1. In general about equimolar amounts of ether and alcohol or an excess of the ether will be used to ensure complete conversion of the alcohol. However, it has been observed that a large excess of the ether is undesirable in that it tends to result in the formation of a precipitate of granular crystals (possibly due to polymerisation of products of decomposition of the ether) and we prefer to employ a mole ratio of ether to alcohol of no greater than about 2: 1.

5

10

15

20

25

30

The separation, recovery and purification of Sevoflurane from product streams containing it is known and any of the known methods may be used in the process of the invention. Such methods will usually include at least one distillation step and will usually include a step of separating and recovering any bis(fluoromethyl)ether present in the product stream. The ether recovered from the product stream can be recycled to the reaction with hexafluoroisopropyl alcohol.

The process can be operated as a batch or continuous process or a combination thereof but is preferably operated as a continuous process with recycle of recovered bis(fluoromethyl)ether.

As stated herinbefore, a prefered embodiment of the invention includes the step of producing the bis(fluoromethyl)ether by reaction of formaldehyde (or a polymeric form of formaldehyde such as paraformaldehyde or trioxane) with hydrogen fluoride. Any of the known methods for production of the bis(fluoromethyl)ether may be employed as the ether formation step of this embodiment of the present invention. The production of bis(fluoromethyl)ether from formaldehyde and hydrogen fluoride is described, for example, in European Patent Publication No. 518,506 and in International Patent Publications No. WO 93/10070, WO 93/12057 and WO 93/22265, the disclosures of which are incorporated herein by reference. In the present invention we especially prefer to employ the ether production process described in International Patent Publication No. WO 93/10070 which comprises reacting formaldehyde with hydrogen fluoride in a reaction-distillation column from which the ether is withdrawn in essentially pure form and in particular essentially free from water.

The invention is illustrated but in no way limited by the following Example.

Example.

5

Bis(fluoromethyl)ether (2g) of purity approximately 99% was mixed with stirring into hexafluoroisopropyl alcohol (4.1g) at room temperature (approximately 25°C) and pressure and sulphuric acid (1ml of 98% acid) was added to the mixture. A slight exotherm was noted. After 10 minutes, the mixture was neutralised by adding sodium hydroxide and then analysed by gas chromatography/mass spectometry.

The composition of the product mixture determined by GC/MS analysis was:-

		%age of mixture
	Sevoflurane	59.7
10	Acetal *	21.6
	Bis(fluoromethyl)ether	17.1
	Methyl formate	1.6
		100
15	•	****

For purposes of comparison the product mixture obtained in Example 19 of WO 93/12057 was determined by GC/MS analysis to be:-

		%age of mixture
	Sevoflurane	4.9
20	Acetal *	1.3
	Bis(fluoromethyl)ether	22.2
	Hexafluoroisopropanol	71.6

25

30

CLAIMS:

5

1. A process for the production of fluoromethylhexafluoroisopropylether which comprises reacting essentially pure bis(fluoromethyl)ether with hexafluoroisopropyl alcohol and recovering the resulting fluoromethylhexafluoroisopropylether from the reaction products.

- 2. A process as claimed in claim 1 which comprises reacting formaldehyde with hydrogen fluoride to produce a reaction mixture containing bis(fluoromethyl)ether, separating essentially pure bis(fluoromethyl)ether from the reaction mixture and reacting the essentially pure bis(fluoromethyl)ether with hexafluoroisopropyl alcohol.
- 3. A process as claimed in claims 1 or 2 in which the essentially pure bis(fluoromethyl)ether is reacted with hexafluoroisopropyl alcohol in the presence of an acid.
 - 4. A process as claimed in any preceding claim in which the molar ratio of bis(fluoromethyl)ether to hexafluoroisopropyl alcohol is 0.5:1 to 2:1.
- 5. A process as claimed in any preceding claim in which the yield of fluoromethyl(hexafluoromethylhexafluoroisopropylether is at least about 55%.

20

25


30

INTERNATIONAL SEARCH REPORT

inte onal Application No
PCT/GB 96/03001

_			,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,
A. CLASS IPC 6	IFICATION OF SUBJECT MATTER C07C41/14 C07C43/12	-	
According	to International Patent Classification (IPC) or to both national class	sification and IPC	
B. FIELDS	SEARCHED		
Minimum d IPC 6	locumentation searched (classification system followed by classification s	stion symbols)	
Documentar	tion searched other than minimum documentation to the extent that	t such documents are included	in the fields searched
Electronic d	late base consulted during the international search (name of data b	ase and, where practical, searc	h terms used)
C. DOCUM	IBNTS CONSIDERED TO BE RELEVANT		
Category '	Citation of document, with indication, where appropriate, of the	relevant passages	Relevant to claim No.
Α .	WO 93 12057 A (ICI) 24 June 1993 cited in the application see examples 1-8,13-19		1-5
Fuet	ner documents are listed in the continuation of box C.	Y Patent family memb	ers are listed in annex.
<u> </u>		X Patent family memb	er are men in amer.
*Special categories of cited documents: 'A' document designing the general state of the art which is not considered to be of particular relevance 'B' earlier document but published on or after the international filing date 'L' document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) 'O' document referring to an oral disclosure, use, exhibition or other means 'P' document published prior to the international filing date but later than the priority date claimed Date of the actual completion of the international search		"I later document published after the international filing date or priority data and not in conflict with the application but cited to understand the principle or theory underlying the invention. "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone. "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. "&" document member of the same patent family Date of mailing of the international search report 0 3, 03, 97	
Name and n	nailing address of the ISA European Patent Office, P.B. 5818 Patentiaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo ni, Fax: (+31-70) 340-3016	Authorized officer Wright, M	

1



Information on patent family members

Inte .onal Application No PCT/GB 96/03001

Patent document cited in search report	Publication date	Patent family member(s)		Publication date	
WO-A-9312057	24-06-93	BR-A-	2954792 9206904 2124934 1074208 0642486 942696 7502037 942183 5504263	19-07-93 21-11-95 24-06-93 14-07-93 15-03-95 08-06-94 02-03-95 10-06-94 02-04-96	